

On the analytic zero divisor conjecture of Linnell

GÁBOR ELEK

Mathematical Institute of the Hungarian Academy of Sciences

P.O. Box 127, H-1364 Budapest, Hungary

elek@renyi.hu

AMS Subject Classifications: 20C07

Keywords: amenable groups, zero divisor conjecture, von Neumann dimension

Abstract. In this note we prove that in the case of finitely generated amenable groups the classical zero divisor conjecture implies the analytic zero divisor conjecture of Linnell.

1 Introduction

Let G be a discrete group. Denote by $\mathbb{C}G$ the complex group algebra of G . The following conjecture is called the zero divisor conjecture.

Conjecture 1 *Let G be a torsion free group. If $0 \neq \alpha \in \mathbb{C}G$ and $0 \neq \beta \in \mathbb{C}G$, then $\alpha\beta \neq 0$.*

Now let

$$L^2(G) = \{ \alpha : G \rightarrow \mathbb{C} \mid \sum_{g \in G} |\alpha(g)|^2 < \infty \}.$$

Then the Hilbert space $L^2(G)$ is a two-sided $\mathbb{C}G$ -module. In [3], Linnell formulated an analytic version of the zero divisor conjecture.

Conjecture 2 *Let G be a torsion free group. If $0 \neq \alpha \in \mathbb{C}G$ and $0 \neq \beta \in L^2(G)$, then $\alpha\beta \neq 0$.*

Since $\mathbb{C}G \subseteq L^2(G)$, the second conjecture implies the first one. The goal of this paper is to prove that for finitely generated amenable groups the two conjectures are actually equivalent.

Theorem 1 *Let G be a finitely generated amenable group. If $0 \neq \alpha \in \mathbb{C}G$, $0 \neq \beta \in L^2(G)$ and $\alpha\beta = 0$, then there exists $0 \neq \gamma \in \mathbb{C}G$ such that $\alpha\gamma = 0$.*

2 Invariant subspaces and the von Neumann dimension

A linear subspace $V \subseteq L^2(G)$ is called an invariant subspace if it is a right $\mathbb{C}G$ -submodule of $L^2(G)$, that is if $\beta \in V$ and $\gamma \in \mathbb{C}G$, then $\beta\gamma \in V$. The von Neumann dimension of an invariant subspace V is a real number defined the following way [4],

$$\dim_G V = \langle P_V 1_e, 1_e \rangle$$

where P_V denotes the orthogonal projection onto the closure of V and $1_e \in L^2(G)$ takes the value 1 on the unit of G and vanishes everywhere else. We list some properties of the von Neumann dimension [4].

1. If $V \neq 0$, then $\dim_G(V) > 0$.
2. $\dim_G(L^2(G)) = 1$.
3. If $V \subseteq W$ are invariant subspaces, then $\dim_G(V) \leq \dim_G(W)$.
4. If $T : L^2(G) \rightarrow L^2(G)$ is a bounded linear transformation and a right $\mathbb{C}G$ -module homomorphism, then

$$\dim_G(\text{Ker } T) + \dim_G(\text{Ran } T) = 1.$$

For an arbitrary linear subspace $W \subseteq L^2(G)$ and a finite subset $A \subseteq G$ one can consider

$$\dim_A(W) = \frac{\sum_{g \in A} \langle P_W 1_g, 1_g \rangle}{|A|},$$

where $1_g \in L^2(G)$ takes the value 1 on g and vanishes everywhere else. [2]. Obviously, $\dim_A(W)$ is a non-negative real number, and if $W \subseteq Z$, then $\dim_A(W) \leq \dim_A(Z)$. If V is an invariant subspace, then for any finite subset $A \subseteq G$, $\dim_A(V) = \dim_G(V)$. Note that if W is a finite dimensional subspace consisting of elements supported on A , then $\dim_A(W) = \frac{\dim_{\mathbb{C}} W}{|A|}$.

3 Approximation of the von Neumann dimension

The goal of this section is to prove a variant of a result of Dodziuk and Mathai [1]. First let us recall the notion of amenability. Let G be a finitely generated group with symmetric generator set $\{g_1, g_2, \dots, g_k\}$. The generators determine a word-metric d on G such a way that the right multiplications by the elements of G are isometries. The group G is called amenable if it has an exhaustion

$$\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \dots, \bigcup_{k=1}^{\infty} \mathcal{F}_k = G$$

by finite subsets such that for any fixed $r \in \mathbb{N}$,

$$\lim_{k \rightarrow \infty} \frac{|\partial_r \mathcal{F}_k|}{|\mathcal{F}_k|} = 0,$$

where $\partial_r \mathcal{F}_k = \{g \in \mathcal{F}_k \mid d(g, G \setminus \mathcal{F}_k) \leq r\}$. Before stating our proposition we list some notations and definitions. Let $\alpha \in \mathbb{C}G$, then $M_\alpha : L^2(G) \rightarrow L^2(G)$ is the left multiplication by α . Then $\text{Ker } M_\alpha$ is an invariant subspace. The width of α , $w(\alpha)$ is defined as $\max_{g \in G, \alpha(g) \neq 0} d(e, g)$, where e is the unit of G . Let $\mathcal{G}_k = \mathcal{F}_k \setminus \partial_{w(\alpha)} \mathcal{F}_k$. Note that if β is supported on $\mathcal{G}_k$, then $\alpha\beta$ will be supported on $\mathcal{F}_k$. We consider the following spaces,

$$V_k = \{\beta \in L^2(G) \mid \text{supp}(\beta) \subseteq \mathcal{F}_k\}$$

$$W_k = \{\beta \in L^2(G) \mid \text{supp}(\beta) \subseteq \mathcal{G}_k\}$$

Proposition 1 $\dim_G(\text{Ker } M_\alpha) = \lim_{k \rightarrow \infty} \dim_{\mathcal{F}_k}(V_k \cap \text{Ker } M_\alpha)$.

Proof: Let $M_\alpha^k : W_k \rightarrow V_k$ the restriction of M_α onto W_k . Then $\text{Ker } M_\alpha^k \subseteq V_k \cap \text{Ker } M_\alpha$, $\text{Ran } M_\alpha^k \subseteq \text{Ran } M_\alpha$ and $\dim_{\mathbb{C}} \text{Ker } M_\alpha^k + \dim_{\mathbb{C}} \text{Ran } M_\alpha^k = |\mathcal{G}_k|$. Therefore,

$$\dim_{\mathcal{F}_k}(\text{Ker } M_\alpha^k) \leq \dim_{\mathcal{F}_k}(V_k \cap \text{Ker } M_\alpha) \leq \dim_G(\text{Ker } M_\alpha) \quad (1)$$

$$\dim_{\mathcal{F}_k}(\text{Ran } M_\alpha^k) \leq \dim_G(\text{Ran } M_\alpha) \quad (2)$$

$$\dim_{\mathcal{F}_k}(\text{Ker } M_\alpha^k) + \dim_{\mathcal{F}_k}(\text{Ran } M_\alpha^k) = \frac{|\mathcal{G}_k|}{|\mathcal{F}_k|} \quad (3)$$

By (3),

$$\lim_{k \rightarrow \infty} (\dim_{\mathcal{F}_k}(\text{Ker } M_\alpha^k) + \dim_{\mathcal{F}_k}(\text{Ran } M_\alpha^k)) = 1 = \dim_G(\text{Ker } M_\alpha) + \dim_G(\text{Ran } M_\alpha) \quad (4)$$

Hence, by (1) and (2), $\lim_{k \rightarrow \infty} \dim_{\mathcal{F}_k} \text{Ker } M_\alpha^k = \dim_G(\text{Ker } M_\alpha)$, thus the Proposition follows. ■

4 The proof of the Theorem

Let $0 \neq \alpha \in \mathbb{C}G$, $0 \neq \beta \in L^2(G)$ and $\alpha\beta = 0$. Then $\text{Ker } M_\alpha \neq 0$, hence $\dim_G(\text{Ker } M_\alpha) > 0$. By the Proposition, for sufficiently large k , there exists $0 \neq \gamma \in V_k \cap \text{Ker } M_\alpha$. Then $\gamma \in \mathbb{C}G$ and $\alpha\gamma = 0$. ■

References

- [1] J. DODZIUK & V. MATHAI, Approximating L^2 -invariants of amenable covering spaces: a combinatorial approach, *Journal of Functional Analysis* **154** (1998) 359-378.
- [2] B. ECKMANN, Approximating L^2 -Betti numbers of an amenable covering by ordinary Betti numbers, *Comment. Math. Helv* **74** (1999) 150-155.
- [3] P.A. LINNELL, Analytic versions of the zero divisor conjecture. *Geometry and cohomology in group theory (Durham 1994)*, London Math. Soc. Lecture Note Series 252, Cambridge University Press (1998) 209-248.
- [4] P. PANSU, Introduction to L^2 -Betti numbers, *American Mathematical Society. Fields Institute Monographs* **4** (1996) 53-86.